

Screening form for renal denervation (RDN)

PATIENT

1 Patients with uncontrolled blood pressure

- Office systolic blood pressure ≥ 140 mmHg AND/OR
- Uncontrolled out of office blood pressure (home or ABPM)

2 Blood pressure lowering medication

- ≥ 3 maximally tolerated antihypertensive medications (ACE-I or ARB + CCB + diuretic as a double or triple combination)
- < 3 with intolerance

3 Patient has expressed interest in renal denervation

- Yes

EXCLUSION CRITERIA (no candidate)

- Known forms of secondary hypertension listed below*
- Pregnancy
- Under 18 years
- Impaired renal function (eGFR < 40 mL/min/1.73 m²)

ADDITIONAL RISK FACTORS (optional)

- Prior stroke
- Transient Ischemic Attack
- Prior Myocardial Infarction
- Coronary Artery Disease
- Heart Failure
- Atrial Fibrillation
- Smoking
- Diabetes mellitus
- Peripheral Vascular Disease

HTN related hospitalizations (admissions) in last 12 months

.....
Emergency room visits related to HTN in the last 12 months

* Primary aldosteronism, pheochromocytoma/paraganglioma, Cushing's syndrome, primary hyperparathyroidism and mineralocorticoid excess syndromes